




2/20


示されるピオチン化 Mab の M3C11 M1E7 M11F1	*	Fン化 Mab の M11F1	$\Omega = \Pi$	sGPC3 = M6B1	アへの結合の M18D4	阻害 (%) M5B9	M10D2	L9G11	エピトープ位置
M3C11	8'96	96.2	12.0	11.2	10.3	9.1	39.9	33.1	
M13B3	71.3	95.7	15.6	9.3	2.7	-1.8	22.3	24.9	ಡ
M1E7	26.0	83.9	14.3	7.7	9.0	6.4	14.8	38.0	
M3B8	1.8	28.4	97.4	15.2	12.6	3.6	29.0	36.0	Q
MIIFI	2.0	3.9	81.8	5.6	-1.5	5.4	24.6	20.9	
M19B11	-4.1	-9.4	3.1	97.8	44.1	31.0	14.2	8.6-	
M6B1	-5.7	-2.7	13.9	85.6	51.9	44.4	6.7	1.1	ပ
M18D4	0.2	0.5	0.3	38.6	89.0	89.4	19.1	6.7	
M5B9	0.1	2.7	5.3	23.2	77.3	77.3	13.4	10.3	
M10D2	2.7.7	9.0	7.8	-5.9	-21.6	-10.3	79.2	7.2	P
L9G11	2.1	-4.8	6.0	10.1	15.7	6.4	1.2	92.2	9








<u>図</u> 4






















細胞傷害活性(%)



13/20

GC33 WB	0	×	×	×	×	×	0	0	0	×
	GC-4 GN S Q Q A T P K D N E I S T F H N L G N V H S P L K	STFHNLGNVHSPLK	GNSQQATPKDNEIS	GNSQQATP	QQATPKDN	TPKDNEIS	ATPKDNEIST	PKDNEISTFH	DNEISTEHNL	EISTFHNLGN
	GC-4	GC-5	9-05	GC-7	8-05	6-05	GC-11	GC-12	GC-13	GC-14

14/20






図15

ELISA	BIACORE		エピトープ	_ر ا		FACS	န		1
kd	-	5	母母	ウエスタン	СНО/				光 强
(1/s		3 8	FI ISA	ブロッティング	human	HepG2	HuH-7	CHO CHO	光概
ر ² 01x	_	(mann)			GPC3				
6.5	_	0.3			83	59	6	-	◁
N.D.	-	N.D.	æ	GC-2	N.D.	N.D.	N.D.		N.D.
42.7	<u> </u>	2.8			28	16	5	٠	×
467.0	Ľ	17.6	1	7.50	12	9	3	٠	×
3.5	L	0.2	0	t >5	9	4	2	,	×
140.0	Ĺ	10.7	อ		29	26	4	,	0
17.1		1.1			39	44	3	٠	0
N.D.	Ľ	N.D.	,	5	37	34	3	٠	0
49.6	_	6.2	ဎ	4	22	20	2	,	0
139.0	Ľ	23.5			25	23	2	,	0
49.6	H	4.7	P		20	29	3	•	0

17/20

		ELISA	H ሕ	エピトープ		FACS	CS	
クローン 田	アインタイプ	EC50 (nM)	競合 ELISA	ウエスタン ブロッティング	CHO/ Eh GPC3	HepG2	HepG2 HuH-7	СНО
GC199	IgG1	0.10	:		15.5	12.2	2.4	ı
GC202	IgG1	0.10	þ	GC-4	0.9	6.0	1.4	ı
GC33	IgG2a	0.24			82.7	52.0	8.4	1
GC179	IgG2b	5.61	¥	6 7 7	7.1	6.5	3.8	J
GC194	lgG1	3.83	Ι	GC-3	5.0	7.9	1.5	

図(




図18

